

CORAIL® PINNACLE®

Your cementless solution, with a **14% lower risk of revision** when compared to all other cementless constructs on the NJR.¹




The CORAIL® Femoral Stem is available both with and without a collar. Proponents of the use of a collared prosthesis claim that it provides advantages in the early stability of the implant, allowing for earlier post-operative weight bearing, protection against subsidence, and a positive dispersion of the vertical forces via the collar into the medial calcar.^{2,3}

The CORAIL collared femoral stems used with PINNACLE have been shown to have a 28% lower risk of revision when compared to all other cementless hips on the NJR.⁴

The NJR has produced a new report analyzing the performance of the CORAIL® PINNACLE® Cementless Construct. This analysis was commissioned by DePuy Synthes, but conducted and validated by the NJR.

The reports details 150,465 CORAIL PINNACLE Cementless implantations (mean age 65.8, 43.7% male). All metal-on-metal was excluded.



Patients receiving a **CORAIL PINNACLE** Cementless Construct are **14% less likely to be revised** when compared to all other cementless constructs on the NJR (HR 0.86 (0.82, 0.90) $p < 0.001$).¹

Please visit www.corailpinnacle.net for more information



DePuy Synthes

PART OF THE **Johnson & Johnson** FAMILY OF COMPANIES

References

1. According to NJR data from 2003 to 2019 where 150,465 CORAIL PINNACLE Cementless implantations were used in primary THR and compared to all other cementless constructs in NJR. Hazard ratio adjusted for age, gender, year cohort and indications. Bespoke Report: Corail - Pinnacle vs. All other Cementless hips. Report can be accessed at <http://www.corailpinnacle.net/supporting-evidence/overview> 2. H. Strom, O. Nilsson, J. Milbrink, et al. The effect of early weight bearing on migration pattern of the uncemented CLS stem in total hip arthroplasty. J Arthroplasty, 22 (2007), p. 1122 3. Demey G, Fary C, Lustig S, Neyret P, Ait si Selmi T. Does a Collar Improve the Immediate Stability of Uncemented Femoral Hip Stems in Total Hip Arthroplasty? A Bilateral Comparative Cadaver Study. J Arthroplasty 26 (2011), No.8, p. 1549 4. According to NJR data from 2003 to 2019 where 81,685 CORAIL Collared femoral stems with PINNACLE cups were used in primary THR and compared to all other cementless hips in NJR. Hazard ratio adjusted for age, gender, year cohort and indications. Bespoke Implant report CORAIL Collared stem PINNACLE cup NJR. Report can be accessed at <http://www.corailpinnacle.net/supporting-evidence/overview> 5. Bespoke Implant report. CORAIL PINNACLE KM_V2 NJR. Report can be accessed at <http://www.corailpinnacle.net/supporting-evidence/overview>

www.jnjmedicaldevices.com

This publication is not intended for distribution in the USA

© DePuy Synthes Joint Reconstruction, a division of Johnson & Johnson Medical Limited. 2020. All rights reserved.
130520-200110 EMEA
DePuy Synthes, Baird House, 4 Lower Gilmore Bank, Edinburgh EH3 9QP